arXiv:1403.4623v1 [math.RA] 18 Mar 2014

IDEMPOTENTS IN NONASSOCIATIVE ALGEBRAS AND EIGENVECTORS OF QUADRATIC OPERATORS

YU. LYUBICH AND A. TSUKERMAN

ABSTRACT. Let F be a field, $\mathrm{char}(F) \neq 2$. Then every finite-dimensional F-algebra has either an idempotent or an absolute nilpotent if and only if over F every polynomial of odd degree has a root in F. This is also necessary and sufficient for existence of eigenvectors for all quadratic operators in finite-dimensional spaces over F.

1. Introduction

Let F be a field, and let A be an *F-algebra*, i.e. a vector space over F endowed by a multiplication that is a bilinear mapping $(x, y) \mapsto xy$ from the Cartesian square $A \times A$ into A. The associativity, the commutativity and the unitality are not assumed in this general definition.

Let $\dim A = n$, $1 \leq n < \infty$, and let $(e_i)_{i=1}^n$ be a basis of A. Then the coordinates of the product $z = xy$ are

$$\zeta_j = \sum_{i,k=1}^{n} \alpha_{ik,j} \xi_i \eta_k, \quad 1 \leq j \leq n, \tag{1.1}$$

where ξ_i, η_k are the coordinates of x and y, respectively, and $\alpha_{ik,j} \in F$ are the uniquely determined coefficients. Every cubic matrix $[\alpha_{ik,j}]$ can be realized in this context but the algebra is commutative if and only if the square matrices $[\alpha_{ik,j}]_{i,k=1}^n$, $1 \leq j \leq n$, are symmetric.

Any algebra A determines a *quadratic operator* $Vx = x^2$ in the underlying vector space . This mapping $A \to A$ is homogeneous of degree 2, i.e. $V(\alpha x) = \alpha^2 Vx$ for all $x \in A$ and all $\alpha \in F$. With $y = x$ in (1.1) we have

$$\zeta_j = \sum_{i,k=1}^{n} \alpha_{ik,j} \xi_i \xi_k, \quad 1 \leq j \leq n, \tag{1.2}$$

where z_j are the coordinates of Vx. If $\mathrm{char}(F) \neq 2$ then the coefficients $\alpha_{ik,j}$ in (1.2) can be symmetrized, thus in this case all quadratic operators come from the commutative algebras.

Similarly to the linear case, a vector $x \neq 0$ is called an *eigenvector* of V if there exists an *eigenvalue* $\lambda \in F$ such that $Vx = \lambda x$, i.e.

$$\sum_{i,k=1}^{n} \alpha_{ik,j} \xi_i \xi_k = \lambda \xi_j, \quad 1 \leq j \leq n. \tag{1.3}$$

[1] *2000 Mathematics Subject Classification:17A01, 12F05*
Keywords: nonassociative algebras, idempotents, nilpotents, quadratic operators, eigenvectors

The eigenvectors of V are just the basis vectors of the one-dimensional subalgebras of the algebra A.

The set $\sigma(V)$ of eigenvalues of V can be called its *spectrum*. However, only two following cases are substantial.

1) $\lambda = 1$. In this case every eigenvector x is a fixed point of V , so $x^2 = x$, i.e. x is an *idempotent* in the generating algebra A.

2) $\lambda = 0$. In this case every eigenvector x is a root of V, so $x^2 = 0$, i.e. x is an *absolute nilpotent*.

Now consider any $\lambda \in \sigma(V)$, and let x be a corrsponding eigenvector. Then there exists $z \in \mathrm{Span}(x) \setminus \{0\}$ which is either an idempotent or an absolute nilpotent in A. Hence, the intersection $\sigma_p(V) = \sigma(V) \cap \{0, 1\}$ is not empty if and only if $\sigma(V) \neq \emptyset$. Moreover, if $\sigma_p(V) = \{0, 1\}$ then $\sigma(V) = F$; if $\sigma_p(V) = \{1\}$ then $\sigma(V) = F \setminus \{0\}$; if $\sigma_p(V) = \{0\}$ then $\sigma(V) = \{0\}$.

We say that a field F is of *class* Λ if every finite-dimensional F-algebra contains either an idempotent or an absolute nilpotent. Equivalently, $F \in \Lambda$ means that in every finite-dimensional vector space over F every quadratic operator has an eigenvector, i.e. its spectrum is not empty.

In [Ly] it was proven that $\mathbb{R} \in \Lambda$. The proof is topological and very short. Let us reproduce it here.

Let $\|.\|$ be an Euclidean norm in $\mathbb{R}^n$, and let $S = \{x : \|x\| = 1\}$. One can assume that $Vx \neq 0$ for all $x \neq 0$. Then $\hat{V}x = Vx/\|Vx\|$ is a smooth mapping $S \to S$. Its topological degree is even since $\hat{V}(-x) = \hat{V}x$. Therefore, $\hat{V}$ is not homotopic to the identity mapping $S \to S$. Hence, there are $x \in S$ and $\tau \in (0, 1)$ such that $(1 - \tau)\hat{V}x + \tau x = 0$. Thus, x is an eigenvector of V.

Note that if *a field F is of class Λ then every its finite (i.e., finite-dimensional) extension Φ is also of class Λ*. Indeed, let A be an l-dimensional Φ-algebra. Then it is an m-dimensional F-algebra with $m = l[\Phi : F] \equiv l \dim_F \Phi$. Every idempotent or absolute nilpotent in F-setting remains as is when F extends to Φ.

In particular, $\mathbb{C} \in \Lambda$ *since* $\mathbb{R} \in \Lambda$.

In the present paper we establish a necessary and a sufficient condition for $F \in \Lambda$. These conditions coincide if $\mathrm{char}(F) \neq 2$.

2. Results

Denote by Γ the class of fields F such that every polynomial of odd degree with coefficients from F has a root in F. For $F = \mathbb{R}$ this property is the only topological ingredient of the Gauss proof (1815) of the "Fundamental Theorem of Algebra".

Theorem 2.1. *Every algebraically closed field F is of class Λ.*

Our Main Theorem is

Theorem 2.2. $\Gamma' \subset \Lambda \subset \Gamma$ *where* $\Gamma' = \Gamma \cap \{F : \mathrm{char}(F) \neq 2\}$.

Corollary 2.3. *If* $\mathrm{char}(F) \neq 2$ *then* $F \in \Lambda$ *if and only if* $F \in \Gamma$.

With the additional condition $\mathrm{char}(F) \neq 2$ Theorem 2.1 follows from Corollary 2.3. Also, the latter yields a "real " counterpart of Theorem 2.1.

Corollary 2.4. *Every formally real and really closed field F is of class Λ.*

Indeed, in this case $F \in \Gamma'$ since $\mathrm{char}(F) = 0$ and $F \in \Gamma$, see e.g. [La], Ch.11.

In several corollaries below we use the inclusion $\Lambda \subset \Gamma$. For instance, *the field* $\mathbb{Q}$ *of rational numbers is not of class* Λ since the polynomial $\alpha^3 - 2$ has no roots in $\mathbb{Q}$. More generally, we have

Corollary 2.5. *Every finite extension* $F \supset \mathbb{Q}$ *is not of class* Λ.

Proof. Assume $F \in \Lambda$. Let Φ be a finite extension of F normal over $\mathbb{Q}$. With $n = [\Phi : \mathbb{Q}]$ let m be any odd number, $m > n$. By Eisenstein's test there exists an irreducible polynomial f of degree m over $\mathbb{Q}$. This f has no roots in Φ, otherwise, all m roots of f belong to Φ, so $m \leq n$. Since f is a polynomial over Φ, we see that $\Phi \notin \Gamma$. Hence, $\Phi \notin \Lambda$, a fortiori, $F \notin \Lambda$, a contradiction. □

The same argument yields

Corollary 2.6. *Let* p *be a prime number, and let* $\mathbb{Q}_p$ *be the field of* p*-adic numbers. Every finite extension* $F \supset \mathbb{Q}_p$ *is not of class* Λ.

Corollary 2.7. *Every finite field* F *is not of class* Λ.

Proof. If $q = \text{card}(F)$ then $\alpha^q - \alpha + 1 = 1 \neq 0$ for all $\alpha \in F$. This polynomial is of odd degree if $\text{char}F \neq 2$. If $\text{char}F = 2$ then such a polynomial is $\alpha^{q+1} - \alpha^2 + 1$. □

Corollary 2.8. *For every field* K *and every* $n \geq 1$ *the field* $F = K(t_1, \cdots, t_n)$ *of rational functions of* n *variables over* K *is not of class* Λ.

Proof. $\alpha^3 - t_1 \neq 0$ for all $\alpha \in F$. □

The class Γ is closely related to the class Δ of fields such that the degrees of all their finite extensions are powers of 2.

Theorem 2.9. $\Gamma' \subset \Delta \subset \Gamma$.

Corollary 2.10. *If* $char(F) \neq 2$ *then* $F \in \Gamma$ *if and only if* $F \in \Delta$.

Thus, if $\text{char}(F) \neq 2$ then each of conditions $F \in \Gamma$ and $F \in \Delta$ is necessary and sufficient for $F \in \Lambda$.

3. Proofs

We start with the following

Lemma 3.1. *A field* F *is of class* Γ *if and only if the degrees of all its nontrivial finite extensions are even.*

Proof. Let $F \notin \Gamma$, so there exists a polynomial f over F of odd degree without roots in F. Then f has an irreducible factor with the same properties. The corresponding extension of F is of odd degree $d > 1$, a contradiction.

Now let $F \in \Gamma$, and let $\Phi \supset F$ be a finite extension of odd degree $d > 1$. For $x \in \Phi \setminus F$ the degree d_x of the extension $F[x]$ is odd as a divisor of d. However, d_x is the degree of the irreducible polynomial f over F such that $f(x) = 0$. This is a contradiction since f has a root in F. □

This Lemma immediately implies $\Delta \subset \Gamma$ that is a part of Theorem 2.9.

Proof of $\Lambda \subset \Gamma$. Let $F \notin \Gamma$. By Lemma 3.1 there exists an extension $\Phi \supset F$ of odd degree $d > 1$. Consider the projection $\pi : \Phi \to \Phi$ onto the one-dimensional subspace F. We introduce in Φ the new multiplication

$$x \circ y = (x - \pi x)(y - \pi y). \tag{3.1}$$

Then Φ becomes a commutative F-algebra. The subspace F is an ideal in Φ since $x \circ y = 0$ for $x \in F$ and all $y \in \Phi$. In the quotient algebra F-algebra $A = \Phi/F$ we consider the equation

$$X \circ X = \lambda X \tag{3.2}$$

with $\lambda \in F$. To conclude that $F \notin \Lambda$ it suffices to show that (3.2) has no solutions $X \neq 0$. Suppose to the contrary. Then there exists $x \in \Phi \setminus F$ such that

$$(x - \pi x)^2 - \lambda x - \mu = 0$$

with a $\mu \in F$. This is a quadratic equation over F since $\pi x \in F$. Its second root is $x' = 2\pi x + \lambda - x \equiv -x \ (\mathrm{mod} F)$, so $x' \notin F$. Hence, the extension $F[x] \supset F$ is of degree 2. On the other hand, $F[x] \subset \Phi$. Therefore, $d = 2[\Phi : F[x]]$. Thus, d is even, a contradiction. □

Now note that *every field* $F \in \Gamma'$ *is perfect*. Indeed, if $\mathrm{char}(F)$ is an odd prime p then $F^p = F$ because of $F \in \Gamma$. Recall that if a field F is perfect then all its normal algebraic extensions are separable, so they are the Galois extensions that allows us to refer to the Galois theory, see e.g. [La], Ch. 8.

Proof of the inclusion $\Gamma' \subset \Delta$. Let $\Phi \supset F$ be a finite extension of F, and let $\Psi \supset \Phi$ be a Galois extension of F. If $[\Phi : F] = 2^i(2k+1)$ then $[\Psi : F] = 2^j(2l+1)$ where $j \geq i$ and $2l+1$ is divisible by $2k+1$. Consider the Galois group $G = \mathrm{Gal}(\Psi/F)$ and its Sylow 2-subgroup H. Let Ω be the subfield of Ψ consisting of the fixed points of H. The index of H in G is $2l+1$, hence $[\Omega : F] = 2l+1$. Since $F \in \Gamma$, we conclude that $l = 0$ by Lemma 3.1. A fortiori, $k = 0$, hence $[\Phi : F] = 2^i$. □

Theorem 2.9 is already proven. As to Theorem 2.2, we have to prove the inclusion $\Gamma' \subset \Lambda$. To this end (and also to prove Theorem 2.1) we return to (1.3) and consider it as a system of n homogeneous equations of second degree

$$g_j(x, \lambda) \equiv \sum_{i,k=1}^{n} \alpha_{ik,j} \xi_i \xi_k - \lambda \xi_j = 0, \quad 1 \leq j \leq n, \quad x = (\xi_1, \cdots, \xi_n) \in F^n, \tag{3.3}$$

with $n+1$ unknowns $\xi_1, \cdots, \xi_n, \lambda$. With $\lambda \in F$ this is a system of equations in the projective space FP^n. The solution $(\mathbf{0}{:}1) \equiv (0 : \cdots : 0 : 1)$ is called *trivial* . In all nontrivial solutions $(x, \lambda) \in FP^n$ the scalar component λ is an eigenvalue of the quadratic operator under consideration. Thus, our aim is to prove the existence of a nontrivial solution in FP^n.

Let us extend FP^n to the projective space $\overline{F}P^n$ over the algebraic closure $\overline{F}$. We call the system (3.3) *generic over* F if the set of its solutions in $\overline{F}P^n$ is finite. Then the number of solutions (i.e., the sum of its multiplicities) is equal to 2^n by the Bezout Theorem. A solution is called *simple* if its multiplicity is equal to 1.

Lemma 3.2. *If the system* (3.3) *is generic over* F *then the trivial solution is simple.*

Proof. The affine space $\overline{F}^n$ can be identified with the projective Zarisky neighborhood of (**0**:1) in $\overline{F}P^n$ consisiting of the points $(x:1)$, $x \in \overline{F}^n$. The corresponding restrictions of the quadratic forms $g_j(x,\lambda)$ are the polynomials $f_j(x) = g_j(x,1)$.

Let R be the local ring of the point $0 \in \overline{F}^n$, and let M be its maximal ideal. In the quotient $\overline{F}$- vector space M/M^2 the images of $(-\xi_j) \in M$, $1 \le j \le n$, constitute a basis. The images of f_j are the same since $f_j + \xi_j \in M^2$. By Nakayama's Lemma the system $(f_j)_{j=1}^n$ generates the ideal M. Hence, the intersection index of the corresponding divisors is 1. □

Let us emphasize that Lemma 3.2 is true irrespective to $\mathrm{char}(F)$.

Proof of Theorem 2.1. . If $F = \overline{F}$ then either the set of solutions in FP^n is infinite or the number of these solutions is $2^n > 1$. By Lemma 3.2 there exists a nontrivial solution of the system (3.3) in FP^n. □

Later on $F \in \Gamma'$. Let the system (3.3) be generic over F. For every solution we fix the projective coordinates and take them from F whensoever the solution belongs to FP^n. Now we consider the Galois extension $\Psi \supset F$ containing all these quantities. By Theorem 2.9 we have $[\Psi : F] = 2^i$ with an $i \ge 1$. Accordingly, the group G=Gal(Ψ/F) is of order 2^i. Since the coefficients of all $g_j(x,\lambda)$ belong to F, the group G naturally acts on the set of solutions preserving their multiplicities. A solution is G-invariant if and only if it is from FP^n.

Let $z = (x,\lambda) \ne 0$ be a solution of a multiplicity m, and let Z be its G-orbit. All points from Z are solutions of the same multiplicity m. The contribution of Z into the number of solutions equals $\hat{m}$ =card$(Z)m$. In this product the second factor coincides with the index of the stabilizer of z. This index is 2^j with some $j \ge 1$ if z is not a fixed point, i.e. if this solution is not from FP^n. Hence, the number of such solutions is even. Since the number of all solutions is 2^n, the number of those solutions which are from FP^n is even. By Lemma 3.2 there exists a nontrivial solution in FP^n. *The generic case in Theorem 2.2 is settled.*

To complete the proof of Theorem 2.2 we show that an "infinitesimal" perturbation of the system (3.3) is generic over an extension of F.

Lemma 3.3. *Let a field $K \supset F$ is endowed by a non-Archimedean valuation, let A be the valuation ring, and let A^0 be its maximal ideal. Then for $1 \le j \le n$ there are some linear forms $\varphi_j(x)$ with coefficients from A and some $\varepsilon_j \in A^0$ such that the system*

$$g_j(x,\lambda) - \varepsilon_j(\varphi_j(x))^2 = 0, \quad 1 \le j \le n, \tag{3.4}$$

is generic over K.

Proof. Inductively on m, $1 \le m \le n$, we construct the variety

$$V_m = \{(x,\lambda) \in \overline{K}P^n : g_j(x,\lambda) - \varepsilon_j(\varphi_j(x))^2 = 0, 1 \le j \le m\}$$

of dimension $\le n - m$. Then $\dim V_n = 0$, thus the system (3.4) is generic over K.

Since $g_1 \ne 0$, we can take $\varepsilon_1 = 0$ to get $\dim V_1 = n - 1$ with any φ_1. Now let $1 \le m < n$, and let $\dim V_m \le n - m$. Consider the decomposition $V_m = \cup_{1 \le i \le r} X_i$ into irreducible components, and choose any point $(x_i, \lambda_i) \in X_i$, $1 \le i \le r$. Let $g_{m+1}(x_i,\lambda_i) = 0$ for $1 \le i \le s$, while $g_{m+1}(x_i,\lambda_i) \ne 0$ for $s+1 \le i \le r$. Since the field K is infinite, there exists a linear form $\varphi_{m+1}(x)$ on $\overline{K}^n$ with coefficients from

A such that $\varphi_{m+1}(x_i) \neq 0$, $1 \leq i \leq r$. With $\varepsilon_{m+1} \in A^0$ different from 0 and from all fractions

$$\frac{g_{m+1}(x_i, \lambda_i)}{(\varphi_{m+1}(x_i))^2}, \quad s+1 \leq i \leq r,$$

the form $g_{m+1}(x, \lambda) - \varepsilon_{m+1}(\varphi_{m+1}(x))^2$ does not vanish on each component of V_m. Therefore, $\dim V_{m+1} < \dim V_m$, hence $\dim V_{m+1} \leq n - m - 1$. □

For our purposes we need to get $K \in \Gamma'$ in Lemma 3.3. The first step in this direction is the extension of F to the field $L = F((t))$ whose nonzero elements are the formal Laurent series

$$a(t) = t^\nu \sum_{k=0}^{\infty} \alpha_k t^k$$

where $\alpha_k \in F$, $\alpha_0 \neq 0$, $\nu = \nu(a) \in \mathbb{Z}$. On L we have the standard non-Archimedean valuation $v_0(a) = \exp(-\nu(a))$, $a \neq 0$. The ground field F is embedded in L as the field of constants, $v_0(\alpha) = 1$ for $\alpha \in F \setminus \{0\}$.

With the distance $v_0(a-b)$ the set L is a complete metric space. The closed unit ball $A_L = \{a \in L : v_0(a) \leq 1\}$, i.e. the set of regular series, is just the valuation ring of the field L. Its unique maximal ideal is the open ball $A_L^0 = \{a : v_0(a) < 1\}$. The residue field $R_L = A_L/A_L^0$ is isomorphic to F.

The valuation v_0 can be extended to a non-Archimedean valuation v of the algebraic closure $\overline{L}$. The corresponding ring is $A_{\overline{L}} = \{x \in \overline{L} : v(x) \leq 1\}$, its maximal ideal is $A_{\overline{L}}^0 = \{x \in \overline{L} : v(x) < 1\}$. Denote by ρ the natural epimorphism from $A_{\overline{L}}$ to the residue field $R_{\overline{L}} = A_{\overline{L}}/A_{\overline{L}}^0$.

Now let $\mathbf{F}$ be the family of fields E such that $L \subset E \subset \overline{L}$, and let $\mathbf{F}$ be partially ordered by inclusion. On every $E \in \mathbf{F}$ we have the valuation $v|E$. The corresponding residue field is isomorphic to $R_E = \mathrm{Im}(\rho|A_E)$ where $A_E = A_{\overline{L}} \cap E$ is the valuation ring of the field E.

Obviosly, if $E_1, E_2 \in \mathbf{F}$ and $E_1 \subset E_2$ then $A_{E_1} \subset A_{E_2}$ and $R_{E_1} \subset R_{E_2}$. In particular, $R_L \subset R_E \subset R_{\overline{L}}$ for all $E \in \mathbf{F}$. By Zorn's Lemma there exists a field $K \in \mathbf{F}$ which is maximal among $E \in \mathbf{F}$ with $R_E = R_L$. Indeed, let $\{E_i\}$ be a linearly ordered subfamily of $\mathbf{F}$, and let all $R_{E_i} = R_L$. Then $\cup E_i$ is a field $E \in \mathbf{F}$, and $A_E = A_{\overline{L}} \cap (\cup E_i) = \cup (A_{\overline{L}} \cap E_i) = \cup A_{E_i}$, hence $R_E = \cup R_{E_i} = R_L$.

Lemma 3.4. $K \in \Gamma'$.

Proof. Since $\mathrm{char}(K) = \mathrm{char}(F) \neq 2$, we actually have to prove that $K \in \Gamma$. Suppose to the contrary. Then by Lemma 3.1 there exists a finite extension $E \supset K$ of odd degree $d > 1$. One can take $E \subset \overline{L}$ since $K \subset \overline{L}$. We have $E \in \mathbf{F}$, but $R_E \neq R_L$ by maximality of K in $\mathbf{F}$. Hence, there exists $x \in E \setminus K$ such that $\rho x \notin R_L$. Let f be an irreducible polynomial over K such that $f(x) = 0$. Its degree d_f is odd since d_f divides d.

Let us extend the initial field L to a field $L_f \subset K$ by joining of all coefficients of f. Obviosly, $L_f \in \mathbf{F}$. The valuation $v|L_f$ is discrete and L_f is complete. Furthermore, $R_{L_f} = R_L$ since $R_L \subset R_{L_f} \subset R_K = R_L$. Hence, R_{L_f} is isomorphic to F, thus it is a perfect field of class Γ.

Now we consider the field $L_f[x] \in \mathbf{F}$. The degree $[L_f[x] : L_f]$ coincides with d_f since the polynomial f is determined and irreducible over L_f. Hence, this degree is odd. The residue field of $L_f[x]$ is a finite extension of R_{L_f}. Its degree δ divides

$[L_f[x] : L_f]$ by Proposition 18 from [La], Ch.12. Hence, δ is odd. On the other hand, $\delta > 1$ by choice of x. By Lemma 3.1 δ is even, a contradiction. □

We also need the following lemma.

Lemma 3.5. *The ring A_K is the direct sum of the subrings F and A_K^0.*

Proof. First, note that $F \subset A_K$ since $v(x) = 1$ for $x \in F$. For the same reason $F \cap A_K^0 = 0$. Now let $x \in A_K$. Then $\rho x \in R_K = R_L$. Hence, $\rho x = \rho a$ where $a \in L$, whence $x - a \in \ker(\rho | A_K) = A_K^0$. In turn, $a = \alpha + \omega$ where $\alpha \in F$ and $\omega \in A_L^0 \subset A_K^0$. As a result, $x = \alpha + (x - a) + \omega \in F + A_K^0$. □

Since Theorem 2.9 is true in the generic case, the system (3.4) has a nontrivial solution $(x, \lambda) \in KP^n$ by Lemmas 3.3 and 3.4. Let $x = (\xi_i)_{i=1}^n \in K^n \setminus \{0\}$, and let $\|x\| = \max_i v(\xi_i)$. Show that $v(\lambda) \le \|x\|$. Indeed, in the opposite case the division on λ^2 in (3.4) yields

$$\zeta_j = h_j(z), \quad 1 \le j \le n, \tag{3.5}$$

where h_j are some quadratic forms with coefficients from A_K and $z = (\zeta_j)_{i=1}^n = (\xi_j/\lambda)_{i=1}^n \in A_K^0 \setminus \{0\}$. However, from (3.5) it follows that $\|z\| \le \|z\|^2$, hence $\|z\| \ge 1$, a contradiction.

Thus, $\max(\|x\|, v(\lambda)) = \|x\|$. Let $\|x\| = v(\xi_1)$ for definiteness. By division on ξ_1 we get a solution $(x_1, \lambda_1) \in KP^n$ with $\|x_1\| = 1$ and $v(\lambda_1) \le 1$. By Lemma 3.5 $(x_1, \lambda_1) = (\hat{x}, \hat{\lambda}) + (y, \omega)$ where the first summand belongs to F^{n+1} and the second one belongs to $(A_K^0)^{n+1}$. In addition, $\hat{x} \ne 0$, otherwise $x_1 = y$, so $\|x_1\| < 1$. Since A_K^0 is an ideal in A_K, the system (3.4) yields $g_j(\hat{x}, \hat{\lambda}) \in F \cap A_K^0$, $1 \le j \le n$. This implies $g_j(\hat{x}, \hat{\lambda}) = 0$, $1 \le j \le n$, by Lemma 3.5 again. *Theorem 2.9 is proven completely.*

4. Some remarks

Remark 4.1. Over any field any power-associative finite-dimensional algebra has an idempotent or a nilpotent of some degree, see [S], Proposition 3.3. In the latter case if $x^r = 0$, $r \ge 2$, but $x^{r-1} \ne 0$, then x^s is an absolute nilpotent for $s = r - [r/2]$. Thus, if we restrict the definition of class Λ to the power-associative algebras then Λ extends to the class of all fields, so our problem disappears. The same happens trivially underthe the restriction to the unital algebras.

Remark 4.2. The infinite-dimensional version of Λ is empty. Indeed, over any field F there are no idempotents neither absolute nilpotents in the algebra of polynomials $f(t)$ such that $f(0) = 0$.

Remark 4.3. The algebra A from the proof of the inclusion $\Lambda \subset \Gamma$ is commutative. Therefore, this inclusion remains valid if in the definition of Λ we consider the commutative algebras only.

Remark 4.4. In the case $\mathrm{char}(F) = 0$ the Corollary 2.1 follows from $\mathbb{C} \in \Lambda$ by the metamathematical Lefshets Principle.

References

[La] S.Lang, Algebra. Addison-Wessley Publ.Co., 1965.

[Ly] Yu.I.Lyubich, The nilpotency of the Engel commutative algebras of dimension $n \leq 4$ over $\mathbb{R}$. Uspechi Mat.Nauk, **32** , no.1 (1977), 195-196 (In Russian).

[S] R.D.Schafer, An Introduction to Nonassociative Algebras. Acad. Press, New York -London, 1966.

Corresponding Author: Yu. Lyubich, Department of Mathematics, Technion, Haifa 32000, Israel

E-mail address: lyubich@tx.technion.ac.il

A. Tsukerman, 30 Port Royal, Foster City, CA 94404